

Yauba MS
Aikhionbare HA
Ogunrinde GO
Bugaje MA

Significant bacteriuria in children with sickle cell anaemia in a Nigerian tertiary hospital

DOI:<http://dx.doi.org/10.4314/njp.v41i4.12>

Accepted: 18th June 2014

Yauba MS (✉)
Aikhionbare HA, Ogunrinde GO
Bugaje MA
Department of Paediatrics,
Ahmadu Bello University Teaching
Hospital,
PMB 06, Shika-Zaria,
Kaduna state, Nigeria.
Email:saadyko@yahoo.co.uk

Abstract Background: Urinary tract infections (UTI) in children with sickle cell anemia (SCA) may result in long term morbidity and mortality due to chronic renal dysfunction.

Objectives: To evaluate the prevalence of significant bacteriuria among children with SCA and to determine their antimicrobial sensitivity patterns of isolates.

Methods: Two hundred and seventy two children with SCA in steady state (n = 185) and in crises (n = 87) aged 6 months to 15 years had their urine samples screened for significant bacteriuria. The urine samples were collected aseptically and incubated aerobically at 37°C for 24 hours. Children whose urine samples yielded $\geq 10^5$ cfu/ml of bacteria on two consecutive cultures were regarded as having significant bacteriuria. The antimicrobial sensitivity pattern of isolates was determined.

Results: Significant bacteriuria

was detected in 22 (8.1%) of the 272 subjects, 156 boys (57.4%) and 116 (42.6%) girls. The prevalence of significant bacteriuria was higher among those in crisis, 18 (20.7%) than among those in steady than, 4 (2.2%) state: $\chi^2 = 27.323$, $p = 0.001$. The most common organism isolated was *Escherichia coli*, 11 (50.0%). The antibiotic with the best sensitivity was ceftriaxone. Most organisms were resistant to the commonly used antibiotics like cotrimoxazole, amoxicillin and ampicillin.

Conclusion: The prevalence of significant bacteriuria was found to be higher in SCA subjects in crisis (20.7%) than among those in steady state (2.2%). The most prevalent urinary pathogens were sensitive to ceftriaxone but resistant to commonly used antibiotics.

Keywords: Sickle Cell Anaemia, Children, Prevalence, Significant Bacteriuria

Introduction

Urinary tract infection (UTI) is a major cause of morbidity in children with sickle cell anemia (SCA). These children have impaired immunological state and are susceptible to recurrent infections including UTI¹. Urinary tract infections in children with SCA, whether symptomatic or asymptomatic, may ultimately lead to chronic kidney disease due to repeated infarctions of the kidney, papillary necrosis, and inability of the kidneys to concentrate urine². Significant bacteriuria is defined as the presence of 10^5 or more organisms per milliliter of midstream urine³. Asymptomatic bacteriuria can also be defined as quantitative growth of bacteria $\geq 10^5$ colony forming units per milliliter of urine of the same organism, from aseptically collected midstream urine specimen, in the absence of symptoms of UTI, usually in repeated urine samples⁴. periodic study of the pattern of the organisms implicated in childhood UTI, and their antibiotic sensitivity of isolates is very important as this

would contribute to reviewing treatment policies. Also, early treatment of identified cases would limit development of chronic kidney disease in this high-risk group of children. This study was undertaken to determine the prevalence of significant bacteriuria in children with SCA in a tertiary hospital. Early detection and management of ASB in SCA children may retard this progression and reduce the morbidity and mortality from CKD.

Subjects and methods

The study was conducted at the Department of Paediatrics, Ahmadu Bello University Teaching Hospital (ABUTH), Zaria, over a period of six months. The study was prospective, descriptive and cross-sectional. The sample population consisted of consecutively selected children with SCA (in steady state and in crisis) aged 6 months to 15 years. Children with SCA who had been

on antibiotics one week preceding enrolment into the study, those with confirmed (or suspected) congenital urogenital anomalies and those who had recent (<1 week) manipulative urogenital procedure (like catheterization and cystoscopy) were excluded from the study. Those whose parents or guardians did not consent and those with HbSC and other forms of sickle cell disease other than SCA were also excluded. Ethical approval was obtained from the ABUTH Research Committee and a written consent obtained from guardians of subjects. Age, sex and other socio-demographic characteristics were recorded.

The urine samples were collected aseptically using standard techniques as described by Anochie *et al.*,⁵ as follows: the subject's external genitalia was cleaned, mid-stream urine was collected, stored in a refrigerator and then submitted to the laboratory within an hour of collection. The urine samples were then incubated aerobically at 37°C for 24 hours within one hour of collection. Children whose urine samples yielded $\geq 10^5$ cfu/ml of bacteria on two consecutive cultures were regarded as having significant bacteriuria. Mixed growths of more than two species in a single urine sample were regarded as contaminants and therefore disregarded. Organisms were identified using standard identification techniques⁶. Antibiotic sensitivity test was also determined using standard methods⁷.

Data analysis

Data was analyzed using Epi Info version 3.5.3 statistical software. Values for continuous variables were expressed as frequency, mean and standard deviation. Chi-square test was used to compare subgroups. P-values less than 0.05 were considered significant.

Results

Socio-demographic characteristics

The ages of the 272 subjects with SCA analyzed ranged from six (6) months to 15 years with a mean age (\pm 1 SD) of 6.4 \pm 3.8 years. Of these 185 (68.0%) were in steady state and 87 (32.0%) were in crisis (Table 1). The mean age of SCA subjects in steady state was 6.8 \pm 3.9 years while that of those in crisis was 5.6 \pm 3.7 years ($p = 0.00$). There were 156 (57.4%) males and 116 (42.6%) females in the ratio of 1.3: 1 with no gender difference ($p = 0.96$). The sex distribution in children in steady state was also not different from that of children presenting in crises ($\chi^2 = 0.084$, $p = 0.39$).

Table 1: Age and gender distribution of SCA subjects in crisis and in steady state

Ages (Years)	In steady state, n (%)		In crisis, n (%)		Total
	Male	Female	Male	Female	
< 5	35 (33.3)	27 (33.8)	26 (51.0)	18 (50.0)	106 (84.1)
5-9	40 (38.1)	35 (43.8)	16 (31.4)	11 (30.6)	102 (71.5)
10-15	30 (28.6)	18 (22.5)	9 (17.6)	7 (19.4)	64 (44.3)
Total	105 (100.0)	80 (100.0)	51 (100.0)	36 (100.0)	272 (100.0)

Prevalence of children with significant bacteriuria

Significant bacteriuria was detected in 22(8.1%) of the 272 subjects with SCA (Table 2). There were 12(54.5%) males and 10(45.5%) females with significant bacteriuria ($p = 0.96$). Among the 22 SCA subjects with significant bacteriuria, 13(59.1%) were less than 5 years, 6(27.3%) were between 5 and 9 years and 3(13.6%) were between 10 and 15 years old. The apparent decrease in significant bacteriuria with increasing age is not significant between the age groups ($p = 0.13$). The prevalence of significant bacteriuria was higher among those in crisis, 18(20.7%), than among those in steady state, 4(2.2%) $-\chi^2 = 27.323$, $p = 0.001$. Fifteen (68.2%) of the 22 subjects with SCA with bacteriuria, were drawn from low social class, 3(13.6%) from the middle class, and 4(18.2%) from the upper class families. Family social class was not associated with prevalence of significant bacteriuria - Fisher exact, $p = 0.19$.

Table 2: Distribution of children with significant bacteriuria by age and clinical status

Age (years)	Children with SB, n (%)		Children without SB, n (%)		Total
	In steady state	In crisis	In steady state	In crisis	
<5	4 (100.0)	9 (50.0)	58 (32.1)	35 (50.7)	106 (39.0)
5-9	0 (0.0)	6 (33.3)	75 (41.4)	21 (30.5)	102 (37.5)
10-15	0 (0.0)	3 (16.7)	48 (26.5)	13 (18.8)	64 (23.5)
Total	4 (100.0)	18 (100.0)	181 (100.0)	69 (100.0)	272 (100.0)

$$\chi^2 = 4.150, df = 2, p = 0.13$$

Bacterial isolates

A total of 22 organisms were isolated from the 22 children with SCA (Table 3). The most frequently isolated organism was *Escherichia coli* (11; 50.0%) followed by *Klebsiella pneumoniae* (5; 22.7%) and *Proteus* species (3; 13.6). The least common isolates were *Staphylococcus aureus* (2; 9.1%) and *Salmonella typhi* (1; 4.6%) Table 3. Antimicrobial sensitivity test revealed that *Escherichia coli*, *Klebsiella pneumoniae* and *Proteus* species were sensitive to gentamycin (95.5%), ciprofloxacin (100%) and ceftriaxone (100%). The organisms were also sensitive to nitrofurantoin (68.2%), cephalexine (61.9%), nalidixic acid (54.5%) and clavulonic acid potentiated amoxicillin (45.5%). The isolates were least sensitive to ampicillin (9.1%), cotrimoxazole (13.6%) and amoxicillin (18.2%).

Table 3: Urinary bacterial isolates among patients with SCA

Organisms	Frequency (%)
<i>Escherichia coli</i>	11 (50.0)
<i>Klebsiella</i> species	5 (22.7)
<i>Proteus</i> species	3 (13.6)
<i>Staphylococcus aureus</i>	2 (9.1)
<i>Salmonella typhi</i>	1 (4.6)
Total	22 (100.0)

Discussion

The overall prevalence of significant bacteriuria in children with SCA in this study was found to be 8.1%. Higher prevalence of significant bacteriuria of 26.7% was reported amongst HbSS patients in Zaria⁸, 26.0% in Maiduguri⁹ all in Nigeria and 10.9% in Jamaica¹⁰. The prevalence observed in this study was also lower than 12.85% reported by Akinbami *et al*¹¹ in Lagos and 21.6% reported by Asinobi *et al*¹² in Ibadan. The lower prevalence of significant bacteriuria obtained in our study could be due to the fact that most of our study population were SCA subjects in steady state, while Zaria⁸, Maiduguri⁹ and Ibadan¹² studies were febrile children with SCA. Differences in the methodology may have contributed to the lower prevalence observed in our study as compared to studies by Akinbami *et al*¹¹ and Cumming *et al*.¹⁰ Whilst our study used clean-catch, mid-stream and supra pubic aspiration urine specimen, Cumming and Akinbami *et al*¹¹ used only clean catch, midstream urine sample for their study. Supra pubic aspiration is more aseptic technique with reduced contamination rate.

However, the prevalence observed in this study was higher than the 6.0% reported by Chuckwu *et al*¹³ in Enugu. The higher prevalence observed in our study as compared to that reported by Chuckwu *et al*¹³ may be contributed by those children with SCA crisis. This study included children with SCA both in crisis and in steady state while Chuckwu *et al*¹³ included only children in steady state.

This study revealed that prevalence of significant bacteriuria did not decrease with increasing age a finding different from that reported by Akuse⁸ and Mava *et al*⁹ where their prevalence decreases with increasing age. The differences in the genetic constitutions among the study populations may explain the difference in the prevalence. Regarding those in crisis, significant bacteriuria was also found to be commoner among the age group less than 5 years than in the older children, a finding similar to that of Akuse⁷ and Mava *et al*⁹. Preponderance of significant bacteriuria in this age group could be explained by the fact of incompletely developed immunity^{8,9}.

Our study revealed that significant bacteriuria was not associated with gender even though it had been shown that bacteriuria is commoner in females than males in children older than one year^{9,12,14}. This could be attributed to the small number of children with confirmed significant bacteriuria in the present study and the varying cultural/religious practices in these environments. In the environment where the current study was conducted male children are not often circumcised before the age of seven years. Early circumcision is the norm in Southern parts of Nigeria and the Western world where Akinola *et al*¹⁴ conducted their study. Lack of circumcision has been associated with increased risk of UTI in boys¹⁵. Significant bacteriuria being not associated with child's socio-economic status contrasted with the finding by Chukwu *et al*¹³.

The current study also showed that the percentage of patients with significant bacteriuria was significantly higher in children with SCA crisis than in those in steady state. The 20.7% prevalence of significant bacteriuria in children with sickle cell crises obtained in this study is in agreement with the 21.6% prevalence reported earlier by Asinobi *et al*¹². On the other hand, the 20.7% prevalence of significant bacteriuria observed in this study in those with SCA crisis was higher than that (10.0%) observed by Tarry *et al*¹⁶ in USA and 25.6% reported by Elbasher and Badu¹⁷ in Saudi Arabia. Differences in the study design and age distribution of study population used might have contributed to the difference in the prevalence. The study by Tarry *et al*¹⁶ was retrospective and included children above 15 years of age while this study included children less than 15 years. This study included only children with SCA while Elbasher and Badu¹⁷ included other haemoglobin phenotypes and their study was retrospective in design, which may explain the difference observed in their study as compared to the present study. Racial variations may also contribute to the differences in the prevalence.

The 2.2% prevalence of significant bacteriuria in steady state SCA subjects obtained in our study was lower than the 6.0% prevalence obtained by Chukwu *et al*¹³ in Enugu and the 5.8% reported by Ajasin and Adegbola¹⁵ in Lagos in children with SCA in steady state. The lower prevalence observed in this study as compared to those of Chukwu *et al*¹³ and Ajasin and Adegbola¹⁸ may be due to the difference in methods of urine collection. Whilst this study used clean-catch, mid-stream and supra pubic aspiration specimen, Chukwu *et al*¹³ and Ajasin and Adegbola¹⁸ used only clean catch, midstream urine sample for their study.

The most frequently isolated organism from the urine specimens was *Escherichia coli*, followed by *Klebsiella* spp, a finding similar to that of other studies^{9,14}. The virulent factors associated with *E.coli*¹⁸ may have contributed to the organisms being the most frequently isolated. *Salmonella typhi* and *Staphylococcus aureus* were the organisms least isolated in our study, a finding similar to that reported by others⁹. The rarity of *Salmonella* UTI in our study as well as these other studies was because *Salmonella typhi* is not a common cause of UTI in children.

This study revealed that most of the pathogens isolated were resistant to cotrimoxazole, ampicillin, cephalixin and amoxicillin but highly sensitive to ceftriaxone, ciprofloxacin, gentamycin, cefuroxime and nitrofurantoin. These findings were similar to those reported by other workers^{12,19}. This may be partly attributed to the high cost of these drugs.

Conclusion

The prevalence of significant bacteriuria was found to be higher in SCA subjects in crisis than in those in

steady state. *Escherichia coli* still remained the commonest pathogen isolated. There was a high degree of resistance of isolated urinary organisms to commonly used antibiotics. It is recommended that routine urine screening for UTI be carried out in children, particularly under-fives, presenting to our hospitals with SCA crisis. Ceftriaxone, cefuroxime or gentamycin should be a drug of choice for empiric treatment of UTI in children with SCA.

Authors' contributions

MS conceived the study, participated in the design and coordination of the study, collected the samples, contributed in the laboratory work, analyzed the data and wrote the final manuscript. HA, GO and MA participated in the design, writing, analyzing and supervision of the study. All contributors approved the final version of the manuscript.

Conflict of interest: None

Funding: None

Acknowledgements

We wish to acknowledge Dr. Adamu Ali Zainab of the General Out-patient Department, ABUTH, Dr. Suleiman Saidu Bashir of the Department of Community Medicine, ABUTH for their immense technical assistance as well as Dr. Ibrahim Adulrasul of the Department of Microbiology, ABUTH for assisting in the laboratory aspect of this study.

References

- Kizito ME, Mworozzi E, Ndugwana C, Serjeant GR. Bacteraemia in homozygous sickle cell disease in Africa: Is pneumococcal prophylaxis justified? *Arch Dis Child*. 2007; 92:21-23
- Chukwu BF, Okafor HU, Ikefuna AN. Socio-demographic factors associated with asymptomatic bacteriuria in children with sickle cell anemia in a tertiary health facility in South eastern, Nigeria. *Italian J Pediatr*. 2011; 3: 1-5.
- Okafor HU, Okoro BA, Ibe BC, Njoku-Obi NU. Prevalence of asymptomatic bacteriuria among nursery school children. *Niger J Paediatr* 1993; 20: 84-88.
- Eke F, Eke NN. Urinary tract infection. In: Azubuike JC, Nkanginieme KEO, editors. Paediatrics and Child Health in a Tropical Region. 1st ed. Owerri: African Education Service; 1999. pp. 326-8.
- Anochie IC, Nkanginieme KEO, Eke FU. The influence of instruction about the methods of urine collection and storage on prevalence of urinary tract infection. *Niger J Paediatr* 2002; 28: 39-42.
- Nester EW, Robert EC, Pearsall NN, Anderson DG, Nester MT. Dynamics of Bacterial Growth. In: Microbiology and Human Perspective. MacGraw-Hall: Boston, 1998: 85-106.
- Cheesbrough M. District laboratory practice in tropical countries. In: Cheesbrough M (ed). Cambridge University Press, United Kingdom, 2000: 124-145.
- Akuse RM. The Pattern of Bacterial Infection and Intercurrent Morbidity in Sickle Cell Disease Patients. A Dissertation Submitted to the National Postgraduate Medical College of Nigeria in Part Fulfillment of the Requirements for the Fellowship of the College 1990: 318-323.
- Mava Y, Ambe JP, Bello M, Watila I, Pius S. Evaluation of the nitrite test in screening for urinary tract infection in febrile children with sickle Cell anaemia in Maiduguri- Nigeria. *Niger Med J* 2011; 52: 45-48.
- Cumming V, Ali S, Forrester T, Roye-Green K, Reid M. Asymptomatic bacteriuria in sickle cell disease: A cross-sectional study. *BMC Infect Dis* 2006;6:46.
- Akinbami AA, Ajibola S, Bode-Shojobi I, Oshinaike O, Adediran A, Ojelabi O *et al.* Prevalence of significant bacteriuria among symptomatic and asymptomatic homozygous sickle cell disease patients in a tertiary hospital in Lagos, Nigeria. *Niger J Clin Pract* 2014; 17: 163 - 167. <http://www.njcponline.com>
- Asinobi AO, Fatunde OJ, Brown BJ, Osinusi K, Fasina NA. Urinary tract infection in febrile children with sickle cell anaemia in Ibadan, Nigeria. *Ann Trop. Paediatr* 2003; 23: 129-134.
- Chukwu BF, Okafor HU, Ikefuna AN. Asymptomatic bacteriuria and sensitivity pattern in children with sickle cell anaemia in a tertiary health centre in Enugu, South East Nigeria. *Ital J Pediatr*. 2011; 37: 45.
- Akinola NO, Stevens SME, Franklin IM, Nash GB, Stuart J. Sub-clinical ischaemic episodes during the steady state of sickle cell anaemia. *J Clin Pathol* 1992; 45: 902-6.
- Herzog L. Urinary tract infections and circumcision. *Am J Dis Child* 1989; 143: 349-350.
- Tarry WF, Dukket JW, McSynder. Urological complications of sickle cell disease in a paediatric population. *J Urol* 1987; 138: 592-594.
- Elbasher AM, Badu GA. Pattern of bacteriuria in patients with sickle cell disease in Qatif Central Hospital. *Saudi Med J* 1991; 12: 121-124.
- Ajasin MA, Adegbola RA. Asymptomatic bacteriuria in children with sickle cell anaemia. *Niger J Paediatr* 1988; 15: 19-25.
- Owa JA. Urinary Tract Infections in Children. In: Azubuike JC, Nkanginieme KEO (editors) Paediatrics and Child health in a Tropical Region. Second ed. Owerri, African Educational Service: 2007: 480-487.